

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
3 July 2003 (03.07.2003)

PCT

(10) International Publication Number  
**WO 03/054179 A1**

(51) International Patent Classification<sup>7</sup>: **C12N 9/12**,  
15/31, 15/52, C07K 14/34, 14/345, C12N 15/77, 1/21,  
C12P 13/08

(21) International Application Number: **PCT/EP02/11561**

(22) International Filing Date: 16 October 2002 (16.10.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
101 62 729.7 20 December 2001 (20.12.2001) DE

(71) Applicant (for all designated States except US): **DE-GUSSA AG** [DE/DE]; Bennigsenplatz 1, 40474 Düsseldorf (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BATHE, Brigitte** [DE/DE]; Twieten 1, 33154 Salzkotten (DE). **HANS, Stephan** [DE/DE]; Wilhelmstrasse 7, 49078 Osnabrück (DE). **KREUTZER, Caroline** [DE/DE]; Poststrasse 16,

49326 Melle (DE). **PFEFFERLE, Walter** [DE/DE]; Jahnstrasse 33, 33790 Halle (DE).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Declarations under Rule 4.17:**

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI,

[Continued on next page]

(54) Title: ALLELES OF THE SIGA GENE FROM CORYNEFORM BACTERIA



(57) Abstract: The invention relates to alleles of the sigA gene from coryneform bacteria which code for sigma factors A and a process for the fermentative preparation of L-lysine using bacteria which contain these alleles.

WO 03/054179 A1



GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

— of inventorship (Rule 4.17(iv)) for US only

**Published:**

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

## Alleles of the sigA gene from coryneform bacteria

### Field of the Invention

The invention provides alleles of the sigA gene from coryneform bacteria which code for variants of sigma factor A and a process for the fermentative preparation of L-lysine using bacteria which contain these alleles.

### Prior art

The amino acid L-lysine is used in human medicine and in the pharmaceuticals industry, in the foodstuffs industry and very particularly in animal nutrition.

It is known that amino acids are prepared by fermentation from strains of coryneform bacteria, in particular *Corynebacterium glutamicum*. Because of their great importance, work is constantly being undertaken to improve the preparation processes. Improvements to the process can relate to fermentation measures, such as, for example, stirring and supply of oxygen, or the composition of the nutrient media, such as, for example, the sugar concentration during the fermentation, or the working up to the product form by, for example, ion exchange chromatography, or the intrinsic output properties of the microorganism itself.

Methods of mutagenesis, selection and mutant selection are used to improve the output properties of these microorganisms. Strains which are resistant to antimetabolites or are auxotrophic for metabolites of regulatory importance and which produce amino acids are obtained in this manner. A known antimetabolite is the lysine analogue S-(2-aminoethyl)-L-cysteine (AEC).

Methods of the recombinant DNA technique have also been employed for some years for improving the strain of *Corynebacterium* strains which produce L-amino acid, by

amplifying individual amino acid biosynthesis genes and investigating the effect on the amino acid production.

The nucleotide sequence of the gene which codes for sigma factor A from *Corynebacterium glutamicum* can be found in  
5 the patent application EP-A-1108790 as sequence no. 2100 and as sequence no. 7065.

The nucleotide sequence is also deposited in the databank of the National Center for Biotechnology Information (NCBI) of the National Library of Medicine (Bethesda, MD, USA)  
10 under Accession Number AX122184 and under Accession Number AX127149.

Object of the invention

The inventors had the object of providing new measures for improved fermentative preparation of L-lysine.

#### 15 Summary of the invention

When L-lysine or lysine are mentioned in the following, not only the bases but also the salts, such as e.g. lysine monohydrochloride or lysine sulfate, are meant by this.

The invention provides replicatable nucleotide sequences  
20 (DNA) which originate from coryneform bacteria, in particular *Corynebacterium glutamicum*, and code for sigma factor A, wherein the associated amino acid sequences in SEQ ID No. 2 contains any proteinogenic amino acid excluding L-alanine at position 414.

25 The invention furthermore provides a replicatable nucleotide sequence (DNA) which originates from coryneform bacteria, in particular *Corynebacterium glutamicum*, and codes for sigma factor A, wherein the associated amino acid sequence contains L-valine at position 414, shown in SEQ ID  
30 No. 4.

The invention furthermore provides a replicatable nucleotide sequence (DNA) which originates from coryneform bacteria, in particular *Corynebacterium glutamicum*, and codes for sigma factor A, the base sequence of which  
5 contains thymine at position 1241, shown in SEQ ID No. 3.

The invention furthermore provides plasmids (vectors) which contain the nucleotide sequences according to the invention and optionally replicate in coryneform bacteria.

The invention furthermore provides coryneform bacteria  
10 which contain the nucleotide sequences according to the invention and in which the nucleotide sequences which code for sigma factor A are optionally present in over-expressed form, wherein the associated amino acid sequences contain another proteinogenic amino acid at position 414 of SEQ ID  
15 No. 2.

Over-expression is understood as meaning an increase in the intracellular concentration or activity of the sigma factors A according to the invention.

By over-expression measures, the activity or concentration  
20 of the corresponding protein is in general increased by at least 10%, 25%, 50%, 75%, 100%, 150%, 200%, 300%, 400% or 500%, up to a maximum of 1000% or 2000%, based on the activity or concentration of the protein in the starting microorganism.

25 Sigma factor A is a transcription factor which mediates the binding of RNA polymerase to specific sites (initiation sites) of the DNA and initiates the start (initiation) of transcription. It participates in the initiation of transcription a large number of genes, for example the  
30 genes *hom*, which codes for homoserine dehydrogenase, *gap*, which codes for glyceraldehyde 3-phosphate dehydrogenase, *fda*, which codes for fructose bisphosphate aldolase, and

pgk, which codes for phosphoglycerate kinase (Pátek et al., Microbiology 143: 1297-1309 (1996)).

To achieve an over-expression, the number of copies of the corresponding genes can be increased, or the promoter and regulation region or the ribosome binding site upstream of the structural gene can be mutated. Expression cassettes which are incorporated upstream of the structural gene act in the same way. By inducible promoters, it is additionally possible to increase the expression in the course of fermentative L-lysine production. The expression is likewise improved by measures to prolong the life of the mRNA. Furthermore, the enzyme activity is also increased by preventing the degradation of the enzyme protein. The genes or gene constructs can either be present in plasmids with a varying number of copies, or can be integrated and amplified in the chromosome. Alternatively, an over-expression of the genes in question can furthermore be achieved by changing the composition of the media and the culture procedure.

Plasmids which are replicated in coryneform bacteria are suitable for increasing the number of copies of the sigma alleles according to the invention. Numerous known plasmid vectors, such as e.g. pZ1 (Menkel et al., Applied and Environmental Microbiology (1989) 64: 549-554), pEKEX1 (Eikmanns et al., Gene 102:93-98 (1991)) or pHS2-1 (Sonnen et al., Gene 107:69-74 (1991)) are based on the cryptic plasmids pHM1519, pBL1 or pGA1. Other plasmid vectors, such as e.g. those based on pCG4 (US-A 4,489,160), or pNG2 (Serwold-Davis et al., FEMS Microbiology Letters 66, 119-124 (1990)), or pAG1 (US-A 5,158,891) can be used in the same manner.

The method of chromosomal gene amplification, such as has been described, for example, by Reinscheid et al. (Applied and Environmental Microbiology 60, 126-132 (1994)) for duplication or amplification of the hom-thrB operon, can

furthermore be used to increase the number of copies. In this method, the complete gene or allele is cloned in a plasmid vector which can replicate in a host (typically *E. coli*), but not in *C. glutamicum*. Possible vectors are, for example, pSUP301 (Simon et al., Bio/Technology 1, 784-791 (1983)), pK18mob or pK19mob (Schäfer et al., Gene 145, 69-73 (1994)), pGEM-T (Promega Corporation, Madison, WI, USA), pCR2.1-TOPO (Shuman, Journal of Biological Chemistry 269:32678-84 (1994); US-A 5,487,993), pCR®Blunt (Invitrogen, Groningen, Holland; Bernard et al., Journal of Molecular Biology, 234: 534-541 (1993)), pEM1 (Schrumpf et al., Journal of Bacteriology 173:4510-4516 (1991)) or pBGS8 (Spratt et al., Gene 41: 337-342 (1986)). The plasmid vector which contains the gene or allele to be amplified is then transferred into the desired strain of *C. glutamicum* by conjugation or transformation. The method of conjugation is described, for example, by Schäfer et al. (Applied and Environmental Microbiology 60, 756-759 (1994)). Methods for transformation are described, for example, by Thierbach et al. (Applied Microbiology and Biotechnology 29, 356-362 (1988)), Dunican and Shivnan (Bio/Technology 7, 1067-1070 (1989)) and Tauch et al. (FEMS Microbiological Letters 123, 343-347 (1994)). After homologous recombination by means of a "cross over" event, the resulting strain contains at least two copies of the gene or allele in question.

The increase in protein concentration is detectable via 1- and 2-dimensional protein gel separation and subsequent optical identification of the protein concentration in the gel with appropriate evaluation software. A common method for preparation of the protein gels in the case of coryneform bacteria and for identification of the proteins is the procedure described by Hermann et al. (Electrophoresis, 22:1712-23 (2001)). The protein concentration can also be analysed by western blot hybridization with an antibody specific for the protein to be detected (Sambrook et al., Molecular cloning: a

laboratory manual. 2<sup>nd</sup> Ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989) and subsequent optical evaluation with appropriate software for determination of the concentration (Lohaus and Meyer (1998) Biospektrum 5:32-39; Lottspeich (1999) Angewandte Chemie 111:2630-2647). The activity of DNA-binding proteins can be measured by means of DNA band shift assays (also called gel retardation) (Wilson et al. (2001) Journal of Bacteriology 183:2151-2155). The effect of DNA-binding proteins on the expression of other genes can be detected by various well-described methods of reporter gene assay (Sambrook et al., Molecular cloning: a laboratory manual. 2<sup>nd</sup> Ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989).

The invention provides replicatable, preferably endogenous nucleotide sequences (DNA) which originate from coryneform bacteria and code for the protein sigma factor A, wherein in the associated amino acid sequences the L-alanine at position 414 of SEQ ID No. 2 is replaced by another proteinogenic amino acid, in particular L-valine, shown in SEQ ID No. 4.

The invention also provides replicatable, preferably endogenous nucleotide sequences (DNA) which originate from coryneform bacteria and code for the protein sigma factor A, the associated base sequence of which contains thymine at position 1241, shown in SEQ ID No. 3.

"Endogenous genes" or "endogenous nucleotide sequences" are understood as meaning the genes or nucleotide sequences present in the population of a species.

The invention also provides vectors (plasmids) which contain the nucleotide sequences mentioned and optionally replicate in coryneform bacteria.

Coryneform bacteria which preferably contain the nucleotide sequence(s) mentioned according to the nucleotide sequences



which code for sigma factor A in an over-expressed form are also claimed.

The invention provides a process for the preparation of L-lysine or feedstuffs additives comprising L-lysine in which  
5 in general the following steps are carried out:

- a) fermentation of coryneform bacteria which contain endogenous nucleotide sequences which code for sigma factor A, wherein in the associated amino acid sequences the L-alanine at position 414 is replaced by  
10 another proteinogenic amino acid, preferably L-valine, the alleles of the endogenous sigA gene are over-expressed under conditions suitable for the formation of the sigA gene product sigma factor A,
- b) concentration of the L-lysine in the fermentation  
15 broth,
- c) isolation of the L-lysine or feedstuffs additive comprising L-lysine from the fermentation broth, optionally
- d) with constituents from the fermentation broth and/or  
20 the biomass (> 0 to 100%).

Proteinogenic amino acids are to be understood as meaning all amino acids which are constituents of proteins or polypeptides. These are, in particular: L-aspartic acid, L-asparagine, L-threonine, L-serine, L-glutamic acid, L-  
25 glutamine, glycine, L-alanine, L-cysteine, L-valine, L-methionine, L-isoleucine, L-leucine, L-tyrosine, L-phenylalanine, L-histidine, L-lysine, L-tryptophan, L-proline and L-arginine.

The wild-type form of the sigA gene is contained in wild-  
30 type strains of coryneform bacteria, in particular of the

genus corynebacterium. It is shown in SEQ ID No. 1. The wild-type protein is shown in SEQ ID No. 2.

Of the genus Corynebacterium, the species Corynebacterium glutamicum known to experts is to be mentioned in particular. Known wild-type strains of the species  
5 Corynebacterium glutamicum are, for example

Corynebacterium glutamicum ATCC13032  
Corynebacterium acetoglutamicum ATCC15806  
Corynebacterium acetoacidophilum ATCC13870  
10 Corynebacterium melassecola ATCC17965  
Corynebacterium thermoaminogenes FERM BP-1539  
Brevibacterium flavum ATCC14067  
Brevibacterium lactofermentum ATCC13869 and  
Brevibacterium divaricatum ATCC14020.

15 Strains with the designation "ATCC" can be obtained from the American Type Culture Collection (Manassas, VA, USA). Strains with the designation "FERM" can be obtained from the National Institute of Advanced Industrial Science and Technology (AIST Tsukuba Central 6, 1-1-1 Higashi, Tsukuba  
20 Ibaraki, Japan). The strain of Corynebacterium thermoaminogenes mentioned (FERM BP-1539) and others (FERM BP-1540, FERM BP-1541 and FERM BP-1542) are described in US-A 5,250,434.

To produce the sigA alleles according to the invention  
25 which code for variants of sigma factor A characterized by an amino acid exchange at position 414 of SEQ ID No. 2, mutagenesis methods described in the prior art are used.

Conventional in vivo mutagenesis methods using mutagenic substances, such as, for example, N-methyl-N'-nitro-N-  
30 nitrosoguanidine, or ultraviolet light can be used for the mutagenesis.

In vitro methods, such as, for example, a treatment with hydroxylamine (Miller, J. H.: A Short Course in Bacterial

Genetics. A Laboratory Manual and Handbook for Escherichia coli and Related Bacteria, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, 1992) or mutagenic oligonucleotides (T. A. Brown: Gentechnologie für  
5 Einsteiger [Genetic Engineering for Beginners], Spektrum Akademischer Verlag, Heidelberg, 1993) or the polymerase chain reaction (PCR) such as is described in the handbook by Newton and Graham (PCR, Spektrum Akademischer Verlag, Heidelberg, 1994) can furthermore be used for the  
10 mutagenesis.

Further instructions on generation of mutations can be found in the prior art and in known textbooks of genetics and molecular biology, such as e.g. the textbook by  
Knippers ("Molekulare Genetik", 6th edition, Georg Thieme  
15 Verlag, Stuttgart, Germany, 1995), that by Winnacker ("Gene und Klone", VCH Verlagsgesellschaft, Weinheim, Germany, 1990) or that by Hagemann ("Allgemeine Genetik", Gustav Fischer Verlag, Stuttgart, 1986).

If in vitro methods are used, the sigA gene described in  
20 the prior art is amplified starting from isolated complete DNA of a wild-type strain with the aid of the polymerase chain reaction, optionally cloned in suitable plasmid vectors, and the DNA is then subjected to the mutagenesis process. Instructions for amplification of DNA sequences  
25 with the aid of the polymerase chain reaction (PCR) can be found by the expert, inter alia, in the handbook by Gait: Oligonucleotide Synthesis: A Practical Approach (IRL Press, Oxford, UK, 1984) and in Newton and Graham: PCR (Spektrum Akademischer Verlag, Heidelberg, Germany, 1994). Suitable  
30 sigA alleles are then selected by the processes described above and investigated.

The invention provides a new sigA allele which codes for a variant of sigma factor A and is shown in SEQ ID No. 3.

The sigA alleles according to the invention can be transferred into suitable strains by the method of gene replacement, such as is described by Schwarzer and Pühler (Bio/Technology 9, 84-87 (1991)) or Peters-Wendisch et al. (Microbiology 144, 915 - 927 (1998)). The corresponding sigA allele is cloned here in a vector which is not replicative for *C. glutamicum*, such as, for example, pK18mobsacB or pK19mobsacB (Jäger et al., Journal of Bacteriology 174: 5462-65 (1992)) or pCR®Blunt (Invitrogen, Groningen, Holland; Bernard et al., Journal of Molecular Biology, 234: 534-541 (1993)) and this is then transferred into the desired host of *C. glutamicum* by transformation or conjugation. After homologous recombination by means of a first "cross-over" event which effects integration and a suitable second "cross-over" event which effects excision in the target gene or in the target sequence, the incorporation of the mutation is achieved.

In addition, it may be advantageous for the production of L-amino acids at the same time to enhance, in particular over-express one or more enzymes of the particular biosynthesis pathway, of glycolysis, of anaplerosis, of the citric acid cycle, of the pentose phosphate cycle, of amino acid export and optionally regulatory proteins, in addition to the use of the sigA allele according to the invention. The use of endogenous genes is in general preferred.

"Endogenous genes" or "endogenous nucleotide sequences" are understood as meaning the genes or nucleotide sequences and alleles thereof present in the population of a species.

The term "enhancement" in this connection describes the increase in the intracellular activity or concentration of one or more enzymes or proteins in a microorganism which are coded by the corresponding DNA, for example by increasing the number of copies of the gene or genes, using a potent promoter or using a gene or allele which codes for

a corresponding enzyme or protein having a high activity, and optionally combining these measures. An increase in the activity of the corresponding enzyme protein can also be effected by a reduced sensitivity to inhibitors.

- 5 By enhancement measures, in particular over-expression, the activity or concentration of the corresponding protein is in general increased by at least 10%, 25%, 50%, 75%, 100%, 150%, 200%, 300%, 400% or 500%, up to a maximum of 1000% or 2000%, based on that of the wild-type protein or the
- 10 activity or concentration of the protein in the starting microorganism.

Thus, for the preparation of L-lysine, in addition to the use of the variants of the sigA gene, at the same time one or more of the endogenous genes chosen from the group

15 consisting of

- the dapA gene which codes for dihydrodipicolinate synthase (EP-B 0 197 335),
- the gap gene which codes for glyceraldehyde 3-phosphate dehydrogenase (Eikmanns (1992), Journal of Bacteriology
- 20 174:6076-6086),
- the eno gene which codes for enolase (DE: 19947791.4),
- the tpi gene which codes for triose phosphate isomerase (Eikmanns (1992), Journal of Bacteriology 174:6076-6086),
- the pgk gene which codes for 3-phosphoglycerate kinase
- 25 (Eikmanns (1992), Journal of Bacteriology 174:6076-6086),
- the zwf gene which codes for glucose 6-phosphate dehydrogenase (JP-A-09224661, EP-A-1108790),
- the pyc gene which codes for pyruvate carboxylase (DE-A-198 31 609; EP-A1108790),

- the mqo gene which codes for malate-quinone oxidoreductase (Molenaar et al., European Journal of Biochemistry 254, 395-403 (1998)),
- 5 • the lysC gene which codes for a feed-back resistant aspartate kinase (Accession No. P26512; EP-B-0387527; EP-A-0699759; WO 00/63388),
- the lyse gene which codes for the lysine export protein (DE-A-195 48 222),
- 10 • the zwal gene which codes for the Zwal protein (DE: 19959328,0; DSM 13115)
- the gnd gene which codes for 6-phosphogluconate dehydrogenase (WO 01/71012),
- 15 • the opca gene which codes for a sub-unit of glucose 6-phosphate dehydrogenase (sequence no. 79 from WO 01/00844; WO 01/04322),

can be enhanced, in particular over-expressed.

The enhancement of 6-phosphogluconate dehydrogenase can also be achieved, inter alia, by amino acid exchanges, such as, for example, by exchange of L-proline for L-serine, L-leucine, L-isoleucine or L-threonine at position 158 of the enzyme protein and/or by exchange of L-serine for L-phenylalanine or L-tyrosine at position 361 of the enzyme protein.

The enhancement of the glucose 6-phosphate dehydrogenase sub-unit can also be achieved, inter alia, by amino acid exchanges, such as, for example, by exchange of L-serine by L-phenylalanine or L-tyrosine at position 312 of the enzyme protein.

It may be furthermore advantageous for the production of L-lysine, in addition to the use of the variants of the sigA

gene, at the same time for one or more of the endogenous genes chosen from the group consisting of

- the pck gene which codes for phosphoenol pyruvate carboxykinase (DE 199 50 409.1, DSM 13047),
- 5 • the pgi gene which codes for glucose 6-phosphate isomerase (US 09/396,478, DSM 12969),
- the poxB gene which codes for pyruvate oxidase (DE:1995 1975.7, DSM 13114),
- 10 • the zwa2 gene which codes for the Zwa2 protein (DE: 19959327,2, DSM 13113),
- the fda gene which codes for fructose 1,6-bisphosphate aldolase (Accession No. X17313; von der Osten et al., Molecular Microbiology 3 (11), 1625-1637 (1989)),
- 15 • the hom gene which codes for homoserine dehydrogenase (EP-A -0131171),
- the leuB gene which codes for isopropyl malate dehydrogenase (Pátek et al., Applied Environmental Microbiology 50:43-47(1989)), Accession No. Y09578),
- 20 • the leuC gene which codes for isopropyl malate dehydratase (Accession No. AX121536, sequence no. 1452 from patent EP1108790, Accession No. AX063983, sequence no. 265 from patent WO0100843),
- the thrB gene which codes for homoserine kinase (Peoples, O.W., et al., Molecular Microbiology 2:63-72(1988)) and
- 25 • the pfkB gene which codes for phosphofructokinase (SEQ ID No. 57 from WO 01/00844)

to be attenuated, in particular for the expression thereof to be reduced.

The term "attenuation" in this connection describes the reduction or elimination of the intracellular activity of one or more enzymes or proteins in a microorganism which are coded by the corresponding DNA, for example by using a weak promoter or using a gene or allele which codes for a corresponding enzyme with a low activity or inactivates the corresponding gene or enzyme or protein, and optionally combining these measures.

By attenuation measures, the activity or concentration of the corresponding protein is in general reduced to 0 to 75%, 0 to 50%, 0 to 25%, 0 to 10% or 0 to 5% of the activity or concentration of the wild-type protein or of the activity or concentration of the protein in the starting microorganism.

The attenuation of isopropyl malate dehydrogenase can also be achieved, inter alia, by amino acid exchanges, such as, for example, by exchange of L-glycine for L-aspartate, L-asparagine or L-glutamate at position 131 of the enzyme protein.

The attenuation of isopropyl malate dehydratase can also be achieved, inter alia, by amino acid exchanges, such as, for example, by exchange of L-arginine for L-serine at position 451 or L-glycine for L-aspartate at position 456 of the enzyme protein or a combination thereof.

The attenuation of homoserine dehydrogenase can also be achieved, inter alia, by amino acid exchanges, such as, for example, by exchange of L-asparagine for L-threonine or L-serine at position 118 or L-leucine for L-proline at position 160 of the enzyme protein or a combination thereof.

The attenuation of phosphofructokinase can also be achieved, inter alia, by amino acid exchanges, such as, for



example, by exchange of L-leucine for L-alanine, L-glycine or L-proline at position 109 of the enzyme protein.

- The invention also provides the microorganisms prepared according to the invention, and these can be cultured
- 5 continuously or discontinuously in the batch process (batch culture) or in the fed batch (feed process) or repeated fed batch process (repetitive feed process) for the purpose of production of L-amino acids. A summary of known culture methods is described in the textbook by Chmiel
- 10 (Bioprozesstechnik 1. Einführung in die Bioverfahrenstechnik (Gustav Fischer Verlag, Stuttgart, 1991)) or in the textbook by Storhas (Bioreaktoren und periphere Einrichtungen (Vieweg Verlag, Braunschweig/Wiesbaden, 1994)).
- 15 The culture medium to be used must meet the requirements of the particular strains in a suitable manner. Descriptions of culture media for various microorganisms are contained in the handbook "Manual of Methods for General Bacteriology" of the American Society for Bacteriology
- 20 (Washington D.C., USA, 1981).

Sugars and carbohydrates, such as e.g. glucose, sucrose, lactose, fructose, maltose, molasses, starch and cellulose, oils and fats, such as, for example, soya oil, sunflower oil, groundnut oil and coconut fat, fatty acids, such as,

25 for example, palmitic acid, stearic acid and linoleic acid, alcohols, such as, for example, glycerol and ethanol, and organic acids, such as, for example, acetic acid, can be used as the source of carbon. These substances can be used individually or as a mixture.

- 30 Organic nitrogen-containing compounds, such as peptones, yeast extract, meat extract, malt extract, corn steep liquor, soya bean flour and urea, or inorganic compounds, such as ammonium sulfate, ammonium chloride, ammonium phosphate, ammonium carbonate and ammonium nitrate, can be

used as the source of nitrogen. The sources of nitrogen can be used individually or as a mixture.

- Phosphoric acid, potassium dihydrogen phosphate or dipotassium hydrogen phosphate or the corresponding sodium-containing salts can be used as the source of phosphorus. The culture medium must furthermore comprise salts of metals, such as, for example, magnesium sulfate or iron sulfate, which are necessary for growth. Finally, essential growth substances, such as amino acids and vitamins, can be employed in addition to the abovementioned substances. Suitable precursors can moreover be added to the culture medium. The starting substances mentioned can be added to the culture in the form of a single batch, or can be fed in during the culture in a suitable manner.
- Basic compounds, such as sodium hydroxide, potassium hydroxide, ammonia or aqueous ammonia, or acid compounds, such as phosphoric acid or sulfuric acid, can be employed in a suitable manner to control the pH of the culture. Antifoams, such as, for example, fatty acid polyglycol esters, can be employed to control the development of foam. Suitable substances having a selective action, such as, for example, antibiotics, can be added to the medium to maintain the stability of plasmids. To maintain aerobic conditions, oxygen or oxygen-containing gas mixtures, such as, for example, air, are introduced into the culture. The temperature of the culture is usually 20°C to 45°C, and preferably 25°C to 40°C. Culturing is continued until a maximum of the desired product has formed. This target is usually reached within 10 hours to 160 hours.
- Methods for the determination of L-amino acids are known from the prior art. The analysis can thus be carried out, for example, as described by Spackman et al. (Analytical Chemistry, 30, (1958), 1190) by anion exchange chromatography with subsequent ninhydrin derivatization, or it can be carried out by reversed phase HPLC, for example

as described by Lindroth et al. (Analytical Chemistry (1979) 51: 1167-1174).

The process according to the invention is used for fermentative preparation of L-lysine.

- 5 The concentration of L-lysine can optionally be adjusted to the desired value by addition of L-lysine.

The present invention is explained in more detail in the following with the aid of embodiment examples.

Example 1

- 10 Amplification and sequencing of the DNA of the sigA allele of strain DM1547

- The *Corynebacterium glutamicum* strain DM1547 was prepared by multiple, non-directed mutagenesis, selection and mutant selection from *C. glutamicum* ATCC13032. The strain is  
15 resistant to the lysine analogue S-(2-aminoethyl)-L-cysteine and methionine-sensitive.

- From the strain DM1547, chromosomal DNA is isolated by the conventional methods (Eikmanns et al., Microbiology 140: 1817 - 1828 (1994)). With the aid of the polymerase chain  
20 reaction, a DNA section which carries the sigA gene or allele is amplified. On the basis of the sequence of the sigA gene known for *C. glutamicum* (sequence no. 2100 and sequence no. 7065 from EP1108970), the following primer oligonucleotides are chosen for the PCR:

- 25 sigA-1 (SEQ ID No. 8):  
5' tgatcggctgaccaactcta 3'

sigA-2 (SEQ ID No. 9):  
5' aaggtctcgaatccgagaac 3'

- The primers shown are synthesized by MWG Biotech  
30 (Ebersberg, Germany) and the PCR reaction is carried out by

the standard PCR method of Innis et al. (PCR protocols. A Guide to Methods and Applications, 1990, Academic Press). The primers allow amplification of a DNA section of approx. 1.89 kb in length, which carries the sigA allele.

- 5 The amplified DNA fragment of approx. 1.89 kb in length which carries the sigA allele of the strain DM1547 is identified by electrophoresis in a 0.8% agarose gel, isolated from the gel and purified by conventional methods (QIAquick Gel Extraction Kit, Qiagen, Hilden).
- 10 The nucleotide sequence of the amplified DNA fragment or PCR product is determined by sequencing by MWG Biotech (Ebersberg, Germany). The sequence of the PCR product is shown in SEQ ID No. 5. The sequence of the coding region is shown again in SEQ ID No. 3. The amino acid sequences of
- 15 the associated sigma factor A protein resulting with the aid of the Patentin program are shown in SEQ ID No. 6 and 4.

At position 1241 of the nucleotide sequence of the coding region of the sigA allele of strain DM1547, that is to say

20 at position 1466 of the nucleotide sequence shown in SEQ ID No. 5, is the base thymine. At the corresponding position of the wild-type gene is the base cytosine (SEQ ID No. 1).

At position 414 of the amino acid sequence of sigma factor A of strain DM1547 is the amino acid valine (SEQ ID No. 6

25 and 4). At the corresponding position of the wild-type protein is the amino acid alanine (SEQ ID No. 2).

The sigA allele, which contains the base thymine at position 1241 of the coding region and accordingly codes for a sigma factor A which contains the amino acid valine

30 at position 414 of the amino acid sequence, is called the sigA\_A414V allele in the following. In the designation "sigA\_A414V", A represents L-alanine, V represents L-valine

and 414 indicates the position of the amino acid exchange (see SEQ ID No. 2 and 4).

### Example 2

Replacement of the sigA wild-type gene of strain DSM5715 by  
5 the sigA\_A414V allele

2.1. Production of a DNA fragment which carries the region of the sigA\_A414V allele on which the mutation A414V is located

From the strain DM1547, chromosomal DNA is isolated by the  
10 conventional methods (Eikmanns et al., Microbiology 140: 1817 - 1828 (1994)). A DNA section which carries the region of the sigA\_A414V allele on which the mutation A414V is located is amplified with the aid of the polymerase chain reaction. On the basis of the sequence of the sigA gene  
15 known for *C. glutamicum* (sequence no. 2100 and sequence no. 7065 from EP-A-1108790), the following primer oligonucleotides are chosen for the PCR such that the mutation A414V is located in the central region of the amplification product:

20 sigA\_XL-A1 (SEQ ID No. 10):  
5' ac gaa ttc-cga cgg cga tga ctt cgt ag 3'

sigA\_XL-A2 (SEQ ID No. 11):  
5' tg gaa ttc-cgt tcc acc tcg ctc cat tc 3'

The primers shown are synthesized by MWG Biotech  
25 (Ebersberg, Germany) and the PCR reaction is carried out by the standard PCR method of Innis et al. (PCR protocols. A guide to methods and applications, 1990, Academic Press). The primers allow amplification of a DNA section approx. 1.69 kb in length which carries a region of the sigA\_A414V  
30 allele (SEQ ID No. 7). The primers moreover contain the sequence for a cleavage site of the restriction

endonuclease EcoRI, which is marked by underlining in the nucleotide sequence shown above.

The amplified DNA fragment of approx. 1.69 kb in length which carries the sigA allele of the strain DM1547 is

5 cleaved with the restriction endonuclease EcoRI, identified by electrophoresis in a 0.8% agarose gel and then isolated from the gel and purified by conventional methods (QIAquick Gel Extraction Kit, Qiagen, Hilden).

## 2.2. Construction of the exchange vector

10 pK18mobsacB\_sigA\_A414V

The approx. 1.68 kb long DNA fragment cleaved with the restriction endonuclease EcoRI, which contains a region of the sigA\_A414V allele which carries the mutation A414V, is incorporated by means of replacement mutagenesis with the aid of the sacB system described by Schäfer et al. (Gene, 15 14, 69-73 (1994)) into the chromosome of the C. glutamicum strain DSM5715. This system enables preparation and selection of allele exchanges which take place by homologous recombination.

20 The mobilizable cloning vector pK18mobsacB is digested with the restriction enzyme EcoRI and the ends are dephosphorylated with alkaline phosphatase (Alkaline Phosphatase, Boehringer Mannheim, Germany). The vector prepared in this way is mixed with the sigA\_A414V fragment of approx. 1.68 kb and the mixture is treated with T4 DNA 25 ligase (Amersham-Pharmacia, Freiburg, Germany).

The E. coli strain S17-1 (Simon et al., Bio/Technologie [Bio/Technology] 1: 784-791, 1993) is then transformed with the ligation batch (Hanahan, In. DNA cloning. A practical 30 approach. Vol.1. ILR-Press, Cold Spring Harbor, New York, 1989). Selection of plasmid-carrying cells is made by plating out the transformation batch on LB agar (Sambrook et al., Molecular Cloning: a laboratory manual. 2<sup>nd</sup> Ed.

Cold Spring Harbor, New York, 1989), which was supplemented with 25 mg/l kanamycin.

Plasmid DNA is isolated from a transformant with the aid of the QIAprep Spin Miniprep Kit from Qiagen and checked by  
5 restriction cleavage with the enzyme BamHI and subsequent agarose gel electrophoresis. The plasmid is called pK18mobsacB\_sigA\_A414V and is shown in figure 1.

### 2.3 Allele exchange

The vector pK18mobsacB\_sigA\_A414V mentioned in example 2.2  
10 is transferred by conjugation by a protocol of Schäfer et al. (Journal of Microbiology 172: 1663-1666 (1990)) into the *C. glutamicum* strain DSM5715. The vector cannot replicate independently in DSM5715 and is retained in the cell only if it is present integrated in the chromosome as  
15 the consequence of a recombination event. Selection of transconjugants, i.e. clones with integrated pK18mobsacB\_sigA\_A414V, is made by plating out the conjugation batch on LB agar (Sambrook et al., Molecular Cloning: a laboratory manual. 2<sup>nd</sup> Ed., Cold Spring Harbor,  
20 New York, 1989), which is supplemented with 15 mg/l kanamycin and 50 mg/l nalidixic acid. Kanamycin-resistant transconjugants are plated out on LB agar plates with 25 mg/l kanamycin and incubated for 24 hours at 33°C. For  
25 selection of mutants in which excision of the plasmid has taken place as a consequence of a second recombination event, the clones are cultured unselectively for 30 hours in LB liquid medium and then plated out on LB agar with 10% sucrose and incubated for 16 hours.

The plasmid pK18mobsacB\_sigA\_A414V, like the starting  
30 plasmid pK18mobsacB, contains, in addition to the kanamycin resistance gene, a copy of the sacB gene which codes for levan sucrase from *Bacillus subtilis*. The expression which can be induced by sucrose leads to the formation of levan sucrase, which catalyses the synthesis of the product

levan, which is toxic to *C. glutamicum*. Only those clones in which the integrated pK18mobsacB\_sigA\_A414V has excised as the consequence of a second recombination event therefore grow on LB agar. Depending on the position of the second recombination event with respect to the mutation site, allele exchange or incorporation of the mutation takes place with the excision, or the original copy remains in the chromosome of the host.

Approximately 40 to 50 colonies are tested for the phenotype "growth in the presence of sucrose" and "non-growth in the presence of kanamycin". In 4 colonies which show the phenotype "growth in the presence of sucrose" and "non-growth in the presence of kanamycin", a region of the sigA gene spanning the mutation A414V is sequenced, starting from the sequencing primer sA\_1 (SEQ ID No. 12), by GATC Biotech AG (Constance, Germany) to demonstrate that the mutation of the sigA\_A414V allele is present in the chromosome. The primer sA\_1 used is synthesized for this by GATC:

sA\_1 (SEQ ID No. 12):  
5' aag ttc tcc acc tac gca ac 3'

A clone which contains the base thymine at position 1241 of the sigA gene and thus has the sigA\_A414V allele was identified in this manner. This clone was called strain DSM5715sigA\_A414V.

### Example 3

#### Preparation of lysine

The *C. glutamicum* strain DSM5715sigA\_A414V obtained in example 2 is cultured in a nutrient medium suitable for the production of lysine and the lysine content in the culture supernatant is determined.



For this, the strain is first incubated on an agar plate for 24 hours at 33°C. Starting from this agar plate culture, a preculture is seeded (10 ml medium in a 100 ml conical flask). The medium MM is used as the medium for the  
5 preculture. The preculture is incubated for 24 hours at 33°C at 240 rpm on a shaking machine. A main culture is seeded from this preculture such that the initial OD (660 nm) of the main culture is 0.1. The Medium MM is also used for the main culture.

10

## Medium MM

CSL	5 g/l
MOPS	20 g/l
Glucose (autoclaved separately)	50 g/l
Salts:	
(NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub>	25 g/l
KH <sub>2</sub> PO <sub>4</sub>	0.1 g/l
MgSO <sub>4</sub> * 7 H <sub>2</sub> O	1.0 g/l
CaCl <sub>2</sub> * 2 H <sub>2</sub> O	10 mg/l
FeSO <sub>4</sub> * 7 H <sub>2</sub> O	10 mg/l
MnSO <sub>4</sub> * H <sub>2</sub> O	5.0mg/l
Biotin (sterile-filtered)	0.3 mg/l
Thiamine * HCl (sterile-filtered)	0.2 mg/l
L-Leucine (sterile-filtered)	0.1 g/l
CaCO <sub>3</sub>	25 g/l

The CSL (corn steep liquor), MOPS (morpholinopropanesulfonic acid) and the salt solution are

brought to pH 7 with aqueous ammonia and autoclaved. The sterile substrate and vitamin solutions, as well as the CaCO<sub>3</sub> autoclaved in the dry state, are then added.

5 Culturing is carried out in a 10 ml volume in a 100 ml conical flask with baffles. Culturing is carried out at 33°C and 80% atmospheric humidity.

After 72 hours, the OD is determined at a measurement wavelength of 660 nm with a Biomek 1000 (Beckmann Instruments GmbH, Munich). The amount of lysine formed is  
10 determined with an amino acid analyzer from Eppendorf-BioTronik (Hamburg, Germany) by ion exchange chromatography and post-column derivatization with ninhydrin detection.

The result of the experiment is shown in table 1.

Tabelle 1

Strain	OD (660 nm)	Lysine HCl g/l
DSM5715	8.2	13.57
DSM5715sigA_A414V	8.0	15.21

15

Brief description of the figure:

Figure 1: Map of the plasmid pK18mobsacB\_sigA\_A414V.

The abbreviations and designations used have the following meaning. The base pair numbers stated are approximate  
20 values obtained in the context of reproducibility of measurements.

Kan: Kanamycin resistance gene

EcoRI: Cleavage site of the restriction enzyme  
EcoRI

BamHI: Cleavage site of the restriction enzyme  
BamHI

'sigA: Cloned DNA fragment containing a 3' terminal  
region of the sigA allele (= sigA\_A414V  
allele) and the downstream region

sacB: sacB gene

RP4-mob: mob region with the replication origin for  
the transfer (oriT)

oriV: Replication origin V


**WHAT IS CLAIMED IS:**

1. Replicable nucleotide sequences (DNA) which originate from coryneform bacteria and code for the protein sigma factor A, wherein in the associated amino acid sequences the L-alanine at position 414 of SEQ ID No. 2 is replaced by another proteinogenic amino acid.
2. Replicable nucleotide sequence (DNA) according to claim 1 which originates from coryneform bacteria and codes for the protein sigma factor A, wherein the associated amino acid sequence contains the L-valine at position 414, shown in SEQ ID No. 4.
3. Replicable nucleotide sequence (DNA) according to claim 1 which originates from coryneform bacteria and codes for the protein sigma factor A, the base sequence of which contains thymine at position 1241, shown in SEQ ID No. 3.
4. Plasmids (vectors) which contain the nucleotide sequences according to claims 1 to 3 and optionally replicate in coryneform bacteria.
5. Coryneform bacteria which contain nucleotide sequences according to claims 1 to 4 and in which the nucleotide sequences which code for sigma factor A are preferably present in over-expressed form.
6. Coryneform bacteria which contain nucleotide sequences which code for sigma factor A, wherein in the associated amino acid sequences the L-alanine at position 414 of SEQ ID No. 2 is replaced by another proteinogenic amino acid.
7. Process for the preparation of L-lysine or feedstuffs additives comprising L-lysine, in which the following steps are carried out (comprising):

- 5           a)    fermentation of coryneform bacteria in which  
              alleles of the endogenous sigA gene are over-  
              expressed under conditions which are suitable for  
              the formation of the sigA gene product sigma  
              factor A; and
- b)    isolation of the L-lysine or the feedstuffs  
                      additive comprising L-lysine from the  
                      fermentation broth, the coryneform bacteria  
                      forming the L-lysine.
- 10   8.    Process according to claim 7, wherein coryneform  
          bacteria which contain an allele of the sigA gene are  
          employed, wherein in the associated amino acid  
          sequences the L-alanine at position 414 of SEQ ID No.  
          2 is replaced by another proteinogenic amino acid.
- 15   9.    Process according to claim 7, wherein microorganisms  
          in which further genes of the biosynthesis pathway of  
          L-lysine are additionally over-expressed are employed.
10.   Process according to claim 7, wherein microorganisms  
                  in which the metabolic pathways which reduce the  
20               formation of L-lysine are at least partly eliminated  
                  are employed.
11.   Process for the preparation of L-lysine or feedstuffs  
                  additives comprising L-lysine, in which the following  
                  steps are carried out:
- 25           a)    fermentation of coryneform bacteria which contain  
                  endogenous nucleotide sequences which code for the  
                  protein sigma factor A, wherein in the associated  
                  amino acid sequences the L-alanine at position 414  
                  is replaced by another proteinogenic amino acid,  
30               preferably L-valine,
- b)    concentration of the L-lysine in the fermentation  
                      broth,

- c) isolation of the L-lysine or feedstuffs additive comprising L-lysine from the fermentation broth, optionally
  - d) with constituents from the fermentation broth and/or the biomass (> 0 to 100%).
- 5

Figure 1: Plasmid pK18mobsacB\_sigA\_A414V



## SEQUENCE LISTING

&lt;110&gt; Degussa AG

&lt;120&gt; Alleles of the sigA gene from coryneform bacteria

&lt;130&gt; 010481 BT

&lt;160&gt; 12

&lt;170&gt; PatentIn version 3.1

&lt;210&gt; 1

&lt;211&gt; 1497

&lt;212&gt; DNA

&lt;213&gt; Corynebacterium glutamicum.

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(1494)

&lt;223&gt; sigA wild-type gene

&lt;400&gt; 1

gtg	gag	agc	agc	atg	gta	gaa	aac	aac	gta	gca	aaa	aag	acg	gtc	gct	48
Met	Glu	Ser	Ser	Met	Val	Glu	Asn	Asn	Val	Ala	Lys	Lys	Thr	Val	Ala	
1				5					10					15		

aaa	aag	acc	gca	cgc	aag	acc	gca	cgc	aaa	gca	gcc	ccg	cgc	gtg	gca	96
Lys	Lys	Thr	Ala	Arg	Lys	Thr	Ala	Arg	Lys	Ala	Ala	Pro	Arg	Val	Ala	
			20					25					30			

acc	cca	ttg	gga	gtc	gca	tct	gag	tct	ccc	att	tcg	gcc	acc	cct	gcg	144
Thr	Pro	Leu	Gly	Val	Ala	Ser	Glu	Ser	Pro	Ile	Ser	Ala	Thr	Pro	Ala	
		35					40					45				

cgc	agc	atc	gat	gga	acc	tca	acc	cct	gtt	gaa	gct	gct	gac	acc	ata	192
Arg	Ser	Ile	Asp	Gly	Thr	Ser	Thr	Pro	Val	Glu	Ala	Ala	Asp	Thr	Ile	
	50					55					60					

gag	acc	acc	gcc	cct	gca	gcg	aag	gct	cct	gcg	gcc	aag	gct	ccc	gct	240
Glu	Thr	Thr	Ala	Pro	Ala	Ala	Lys	Ala	Pro	Ala	Ala	Lys	Ala	Pro	Ala	
65					70					75				80		

aaa	aag	gtt	gcc	aag	aag	aca	gct	cgc	aag	gca	cct	gcg	aaa	aag	act	288
Lys	Lys	Val	Ala	Lys	Lys	Thr	Ala	Arg	Lys	Ala	Pro	Ala	Lys	Lys	Thr	
			85						90					95		

gtc	gcc	aag	aaa	gcc	aca	acc	gcc	aag	gct	gca	cct	gca	act	gcc	aag	336
Val	Ala	Lys	Lys	Ala	Thr	Thr	Ala	Lys	Ala	Ala	Pro	Ala	Thr	Ala	Lys	
			100					105					110			

gac	gaa	aac	gca	cct	gtt	gat	gac	gac	gag	gag	aac	ctc	gct	cag	gat	384
Asp	Glu	Asn	Ala	Pro	Val	Asp	Asp	Asp	Glu	Glu	Asn	Leu	Ala	Gln	Asp	
		115					120					125				

gaa	cag	gac	ttc	gac	ggc	gat	gac	ttc	gta	gac	ggc	atc	gaa	gac	gaa	432
Glu	Gln	Asp	Phe	Asp	Gly	Asp	Asp	Phe	Val	Asp	Gly	Ile	Glu	Asp	Glu	
		130				135					140					



gaa gat gaa gac ggc gtc gaa gcc ctc ggt gaa gaa agc gaa gac gac Glu Asp Glu Asp Gly Val Glu Ala Leu Gly Glu Glu Ser Glu Asp Asp 145 150 155 160	480
gaa gag gac ggc tca tcc gtt tgg gat gaa gac gaa tcc gca acc ctg Glu Glu Asp Gly Ser Ser Val Trp Asp Glu Asp Glu Ser Ala Thr Leu 165 170 175	528
cgt cag gca cgt aaa gat gcc gag ctc acc gct tcc gcc gac tct gtt Arg Gln Ala Arg Lys Asp Ala Glu Leu Thr Ala Ser Ala Asp Ser Val 180 185 190	576
cgc gct tac ctg aag caa atc ggt aaa gtt gcc ctg ctg aac gct gaa Arg Ala Tyr Leu Lys Gln Ile Gly Lys Val Ala Leu Leu Asn Ala Glu 195 200 205	624
cag gaa gtc tcc ctg gca aag cgc atc gaa gca ggc ctt tac gcc acc Gln Glu Val Ser Leu Ala Lys Arg Ile Glu Ala Gly Leu Tyr Ala Thr 210 215 220	672
cac cgc atg gag gaa atg gaa gaa gct ttc gca gcc ggt gac aag gac His Arg Met Glu Glu Met Glu Glu Ala Phe Ala Ala Gly Asp Lys Asp 225 230 235 240	720
gcg aaa ctc acc cca gcc gtc aag cgt gac ctc cgc gcc atc gct cgt Ala Lys Leu Thr Pro Ala Val Lys Arg Asp Leu Arg Ala Ile Ala Arg 245 250 255	768
gac ggc cgc aag gcg aaa aac cac ctc ctg gaa gcc aac ctt cgt ctg Asp Gly Arg Lys Ala Lys Asn His Leu Leu Glu Ala Asn Leu Arg Leu 260 265 270	816
gtt gtc tcc ctg gca aag cgc tac acc ggc cgt ggc atg gca ttc ctg Val Val Ser Leu Ala Lys Arg Tyr Thr Gly Arg Gly Met Ala Phe Leu 275 280 285	864
gac ctc atc cag gaa ggc aac ctc ggt ctg att cgt gcc gta gag aag Asp Leu Ile Gln Glu Gly Asn Leu Gly Leu Ile Arg Ala Val Glu Lys 290 295 300	912
ttc gac tac tcc aag ggc tac aag ttc tcc acc tac gca acc tgg tgg Phe Asp Tyr Ser Lys Gly Tyr Lys Phe Ser Thr Tyr Ala Thr Trp Trp 305 310 315 320	960
atc cgt cag gca atc acc cgc gcc atg gcc gac caa gca cga acc atc Ile Arg Gln Ala Ile Thr Arg Ala Met Ala Asp Gln Ala Arg Thr Ile 325 330 335	1008
cgt atc cca gtc cac atg gtt gaa gtg atc aac aaa ctt ggt cgc atc Arg Ile Pro Val His Met Val Glu Val Ile Asn Lys Leu Gly Arg Ile 340 345 350	1056
caa cgt gaa ctc ctt cag gaa ctc ggc cgc gaa cca acc cca cag gaa Gln Arg Glu Leu Leu Gln Glu Leu Gly Arg Glu Pro Thr Pro Gln Glu 355 360 365	1104
ctg tcc aaa gaa atg gac atc tcc gag gaa aag gta ctg gaa atc cag Leu Ser Lys Glu Met Asp Ile Ser Glu Glu Lys Val Leu Glu Ile Gln 370 375 380	1152

cag tac gcc cgc gaa cca atc tcc ctg gac caa acc atc ggc gac gaa 1200  
 Gln Tyr Ala Arg Glu Pro Ile Ser Leu Asp Gln Thr Ile Gly Asp Glu  
 385 390 395 400  
 ggc gac agc cag ctc ggc gac ttc atc gaa gac tcc gaa gcc gtc gtc 1248  
 Gly Asp Ser Gln Leu Gly Asp Phe Ile Glu Asp Ser Glu Ala Val Val  
 405 410 415  
 gca gtc gac gcc gtc tca ttc acc ctg ctg caa gac cag cta cag gac 1296  
 Ala Val Asp Ala Val Ser Phe Thr Leu Leu Gln Asp Gln Leu Gln Asp  
 420 425 430  
 gtc cta gag acc ctc tcc gaa cgt gaa gcc ggc gtg gtt aaa ctc cgc 1344  
 Val Leu Glu Thr Leu Ser Glu Arg Glu Ala Gly Val Val Lys Leu Arg  
 435 440 445  
 ttc gga ctc acc gac gga atg cca cgc act tta gac gaa atc ggc caa 1392  
 Phe Gly Leu Thr Asp Gly Met Pro Arg Thr Leu Asp Glu Ile Gly Gln  
 450 455 460  
 gtt tac ggt gtc acc cgt gag cgc atc cgc cag att gag tcc aag acc 1440  
 Val Tyr Gly Val Thr Arg Glu Arg Ile Arg Gln Ile Glu Ser Lys Thr  
 465 470 475 480  
 atg tct aag ctg cgc cac cca tca cgc tcc cag gtc ctt cgc gac tac 1488  
 Met Ser Lys Leu Arg His Pro Ser Arg Ser Gln Val Leu Arg Asp Tyr  
 485 490 495  
 ctg gac taa 1497  
 Leu Asp

<210> 2  
 <211> 498  
 <212> PRT  
 <213> Corynebacterium glutamicum

<400> 2  
 Met Glu Ser Ser Met Val Glu Asn Asn Val Ala Lys Lys Thr Val Ala  
 1 5 10 15  
 Lys Lys Thr Ala Arg Lys Thr Ala Arg Lys Ala Ala Pro Arg Val Ala  
 20 25 30  
 Thr Pro Leu Gly Val Ala Ser Glu Ser Pro Ile Ser Ala Thr Pro Ala  
 35 40 45  
 Arg Ser Ile Asp Gly Thr Ser Thr Pro Val Glu Ala Ala Asp Thr Ile  
 50 55 60  
 Glu Thr Thr Ala Pro Ala Ala Lys Ala Pro Ala Ala Lys Ala Pro Ala  
 65 70 75 80  
 Lys Lys Val Ala Lys Lys Thr Ala Arg Lys Ala Pro Ala Lys Lys Thr  
 85 90 95  
 Val Ala Lys Lys Ala Thr Thr Ala Lys Ala Ala Pro Ala Thr Ala Lys  
 100 105 110  
 Asp Glu Asn Ala Pro Val Asp Asp Asp Glu Glu Asn Leu Ala Gln Asp  
 115 120 125

Glu Gln Asp Phe Asp Gly Asp Asp Phe Val Asp Gly Ile Glu Asp Glu  
 130 135 140  
 Glu Asp Glu Asp Gly Val Glu Ala Leu Gly Glu Glu Ser Glu Asp Asp  
 145 150 155 160  
 Glu Glu Asp Gly Ser Ser Val Trp Asp Glu Asp Glu Ser Ala Thr Leu  
 165 170 175  
 Arg Gln Ala Arg Lys Asp Ala Glu Leu Thr Ala Ser Ala Asp Ser Val  
 180 185 190  
 Arg Ala Tyr Leu Lys Gln Ile Gly Lys Val Ala Leu Leu Asn Ala Glu  
 195 200 205  
 Gln Glu Val Ser Leu Ala Lys Arg Ile Glu Ala Gly Leu Tyr Ala Thr  
 210 215 220  
 His Arg Met Glu Glu Met Glu Glu Ala Phe Ala Ala Gly Asp Lys Asp  
 225 230 235 240  
 Ala Lys Leu Thr Pro Ala Val Lys Arg Asp Leu Arg Ala Ile Ala Arg  
 245 250 255  
 Asp Gly Arg Lys Ala Lys Asn His Leu Leu Glu Ala Asn Leu Arg Leu  
 260 265 270  
 Val Val Ser Leu Ala Lys Arg Tyr Thr Gly Arg Gly Met Ala Phe Leu  
 275 280 285  
 Asp Leu Ile Gln Glu Gly Asn Leu Gly Leu Ile Arg Ala Val Glu Lys  
 290 295 300  
 Phe Asp Tyr Ser Lys Gly Tyr Lys Phe Ser Thr Tyr Ala Thr Trp Trp  
 305 310 315 320  
 Ile Arg Gln Ala Ile Thr Arg Ala Met Ala Asp Gln Ala Arg Thr Ile  
 325 330 335  
 Arg Ile Pro Val His Met Val Glu Val Ile Asn Lys Leu Gly Arg Ile  
 340 345 350  
 Gln Arg Glu Leu Leu Gln Glu Leu Gly Arg Glu Pro Thr Pro Gln Glu  
 355 360 365  
 Leu Ser Lys Glu Met Asp Ile Ser Glu Glu Lys Val Leu Glu Ile Gln  
 370 375 380  
 Gln Tyr Ala Arg Glu Pro Ile Ser Leu Asp Gln Thr Ile Gly Asp Glu  
 385 390 395 400  
 Gly Asp Ser Gln Leu Gly Asp Phe Ile Glu Asp Ser Glu Ala Val Val  
 405 410 415  
 Ala Val Asp Ala Val Ser Phe Thr Leu Leu Gln Asp Gln Leu Gln Asp  
 420 425 430  
 Val Leu Glu Thr Leu Ser Glu Arg Glu Ala Gly Val Val Lys Leu Arg  
 435 440 445  
 Phe Gly Leu Thr Asp Gly Met Pro Arg Thr Leu Asp Glu Ile Gly Gln  
 450 455 460

Val Tyr Gly Val Thr Arg Glu Arg Ile Arg Gln Ile Glu Ser Lys Thr  
465 470 475 480

Met Ser Lys Leu Arg His Pro Ser Arg Ser Gln Val Leu Arg Asp Tyr  
485 490 495

Leu Asp

<210> 3  
<211> 1497  
<212> DNA  
<213> Corynebacterium glutamicum

<220>  
<221> CDS  
<222> (1)..(1494)  
<223> sigA allele

<220>  
<221> mutation  
<222> (1241)..(1241)  
<223> Exchange of cytosine for thymine

<400> 3  
gtg gag agc agc atg gta gaa aac aac gta gca aaa aag acg gtc gct 48  
Met Glu Ser Ser Met Val Glu Asn Asn Val Ala Lys Lys Thr Val Ala  
1 5 10 15  
aaa aag acc gca cgc aag acc gca cgc aaa gca gcc ccg cgc gtg gca 96  
Lys Lys Thr Ala Arg Lys Thr Ala Arg Lys Ala Ala Pro Arg Val Ala  
20 25 30  
acc cca ttg gga gtc gca tct gag tct ccc att tcg gcc acc cct gcg 144  
Thr Pro Leu Gly Val Ala Ser Glu Ser Pro Ile Ser Ala Thr Pro Ala  
35 40 45  
cgc agc atc gat gga acc tca acc cct gtt gaa gct gct gac acc ata 192  
Arg Ser Ile Asp Gly Thr Ser Thr Pro Val Glu Ala Ala Asp Thr Ile  
50 55 60  
gag acc acc gcc cct gca gcg aag gct cct gcg gcc aag gct ccc gct 240  
Glu Thr Thr Ala Pro Ala Ala Lys Ala Pro Ala Ala Lys Ala Pro Ala  
65 70 75 80  
aaa aag gtt gcc aag aag aca gct cgc aag gca cct gcg aaa aag act 288  
Lys Lys Val Ala Lys Lys Thr Ala Arg Lys Ala Pro Ala Lys Lys Thr  
85 90 95  
gtc gcc aag aaa gcc aca acc gcc aag gct gca cct gca act gcc aag 336  
Val Ala Lys Lys Ala Thr Thr Ala Lys Ala Ala Pro Ala Thr Ala Lys  
100 105 110  
gac gaa aac gca cct gtt gat gac gac gag gag aac ctc gct cag gat 384  
Asp Glu Asn Ala Pro Val Asp Asp Asp Glu Glu Asn Leu Ala Gln Asp  
115 120 125  
gaa cag gac ttc gac ggc gat gac ttc gta gac ggc atc gaa gac gaa 432  
Glu Gln Asp Phe Asp Gly Asp Asp Phe Val Asp Gly Ile Glu Asp Glu  
130 135 140

gaa gat gaa gac ggc gtc gaa gcc ctc ggt gaa gaa agc gaa gac gac Glu Asp Glu Asp Gly Val Glu Ala Leu Gly Glu Glu Ser Glu Asp Asp 145 150 155 160	480
gaa gag gac ggc tca tcc gtt tgg gat gaa gac gaa tcc gca acc ctg Glu Glu Asp Gly Ser Ser Val Trp Asp Glu Asp Glu Ser Ala Thr Leu 165 170 175	528
cgt cag gca cgt aaa gat gcc gag ctc acc gct tcc gcc gac tct gtt Arg Gln Ala Arg Lys Asp Ala Glu Leu Thr Ala Ser Ala Asp Ser Val 180 185 190	576
cgc gct tac ctg aag caa atc ggt aaa gtt gcc ctg ctg aac gct gaa Arg Ala Tyr Leu Lys Gln Ile Gly Lys Val Ala Leu Leu Asn Ala Glu 195 200 205	624
cag gaa gtc tcc ctg gca aag cgc atc gaa gca ggc ctt tac gcc acc Gln Glu Val Ser Leu Ala Lys Arg Ile Glu Ala Gly Leu Tyr Ala Thr 210 215 220	672
cac cgc atg gag gaa atg gaa gaa gct ttc gca gcc ggt gac aag gac His Arg Met Glu Glu Met Glu Glu Ala Phe Ala Ala Gly Asp Lys Asp 225 230 235 240	720
gcg aaa ctc acc cca gcc gtc aag cgt gac ctc cgc gcc atc gct cgt Ala Lys Leu Thr Pro Ala Val Lys Arg Asp Leu Arg Ala Ile Ala Arg 245 250 255	768
gac ggc cgc aag gcg aaa aac cac ctc ctg gaa gcc aac ctt cgt ctg Asp Gly Arg Lys Ala Lys Asn His Leu Leu Glu Ala Asn Leu Arg Leu 260 265 270	816
gtt gtc tcc ctg gca aag cgc tac acc ggc cgt ggc atg gca ttc ctg Val Val Ser Leu Ala Lys Arg Tyr Thr Gly Arg Gly Met Ala Phe Leu 275 280 285	864
gac ctc atc cag gaa ggc aac ctc ggt ctg att cgt gcc gta gag aag Asp Leu Ile Gln Glu Gly Asn Leu Gly Leu Ile Arg Ala Val Glu Lys 290 295 300	912
ttc gac tac tcc aag ggc tac aag ttc tcc acc tac gca acc tgg tgg Phe Asp Tyr Ser Lys Gly Tyr Lys Phe Ser Thr Tyr Ala Thr Trp Trp 305 310 315 320	960
atc cgt cag gca atc acc cgc gcc atg gcc gac caa gca cga acc atc Ile Arg Gln Ala Ile Thr Arg Ala Met Ala Asp Gln Ala Arg Thr Ile 325 330 335	1008
cgt atc cca gtc cac atg gtt gaa gtg atc aac aaa ctt ggt cgc atc Arg Ile Pro Val His Met Val Glu Val Ile Asn Lys Leu Gly Arg Ile 340 345 350	1056
caa cgt gaa ctc ctt cag gaa ctc gcc cgc gaa cca acc cca cag gaa Gln Arg Glu Leu Leu Gln Glu Leu Gly Arg Glu Pro Thr Pro Gln Glu 355 360 365	1104
ctg tcc aaa gaa atg gac atc tcc gag gaa aag gta ctg gaa atc cag Leu Ser Lys Glu Met Asp Ile Ser Glu Glu Lys Val Leu Glu Ile Gln 370 375 380	1152

cag tac gcc cgc gaa cca atc tcc ctg gac caa acc atc ggc gac gaa 1200  
 Gln Tyr Ala Arg Glu Pro Ile Ser Leu Asp Gln Thr Ile Gly Asp Glu  
 385 390 395 400  
 ggc gac agc cag ctc ggc gac ttc atc gaa gac tcc gaa gtc gtc gtc 1248  
 Gly Asp Ser Gln Leu Gly Asp Phe Ile Glu Asp Ser Glu Val Val Val  
 405 410 415  
 gca gtc gac gcc gtc tca ttc acc ctg ctg caa gac cag cta cag gac 1296  
 Ala Val Asp Ala Val Ser Phe Thr Leu Leu Gln Asp Gln Leu Gln Asp  
 420 425 430  
 gtc cta gag acc ctc tcc gaa cgt gaa gcc ggc gtg gtt aaa ctc cgc 1344  
 Val Leu Glu Thr Leu Ser Glu Arg Glu Ala Gly Val Val Lys Leu Arg  
 435 440 445  
 ttc gga ctc acc gac gga atg cca cgc act tta gac gaa atc ggc caa 1392  
 Phe Gly Leu Thr Asp Gly Met Pro Arg Thr Leu Asp Glu Ile Gly Gln  
 450 455 460  
 gtt tac ggt gtc acc cgt gag cgc atc cgc cag att gag tcc aag acc 1440  
 Val Tyr Gly Val Thr Arg Glu Arg Ile Arg Gln Ile Glu Ser Lys Thr  
 465 470 475 480  
 atg tct aag ctg cgc cac cca tca cgc tcc cag gtc ctt cgc gac tac 1488  
 Met Ser Lys Leu Arg His Pro Ser Arg Ser Gln Val Leu Arg Asp Tyr  
 485 490 495  
 ctg gac taa 1497  
 Leu Asp

<210> 4  
 <211> 498  
 <212> PRT  
 <213> *Corynebacterium glutamicum*

<400> 4  
 Met Glu Ser Ser Met Val Glu Asn Asn Val Ala Lys Lys Thr Val Ala  
 1 5 10 15  
 Lys Lys Thr Ala Arg Lys Thr Ala Arg Lys Ala Ala Pro Arg Val Ala  
 20 25 30  
 Thr Pro Leu Gly Val Ala Ser Glu Ser Pro Ile Ser Ala Thr Pro Ala  
 35 40 45  
 Arg Ser Ile Asp Gly Thr Ser Thr Pro Val Glu Ala Ala Asp Thr Ile  
 50 55 60  
 Glu Thr Thr Ala Pro Ala Ala Lys Ala Pro Ala Ala Lys Ala Pro Ala  
 65 70 75 80  
 Lys Lys Val Ala Lys Lys Thr Ala Arg Lys Ala Pro Ala Lys Lys Thr  
 85 90 95  
 Val Ala Lys Lys Ala Thr Thr Ala Lys Ala Ala Pro Ala Thr Ala Lys  
 100 105 110  
 Asp Glu Asn Ala Pro Val Asp Asp Glu Glu Asn Leu Ala Gln Asp  
 115 120 125

Glu Gln Asp Phe Asp Gly Asp Asp Phe Val Asp Gly Ile Glu Asp Glu  
 130 135 140  
 Glu Asp Glu Asp Gly Val Glu Ala Leu Gly Glu Glu Ser Glu Asp Asp  
 145 150 155 160  
 Glu Glu Asp Gly Ser Ser Val Trp Asp Glu Asp Glu Ser Ala Thr Leu  
 165 170 175  
 Arg Gln Ala Arg Lys Asp Ala Glu Leu Thr Ala Ser Ala Asp Ser Val  
 180 185 190  
 Arg Ala Tyr Leu Lys Gln Ile Gly Lys Val Ala Leu Leu Asn Ala Glu  
 195 200 205  
 Gln Glu Val Ser Leu Ala Lys Arg Ile Glu Ala Gly Leu Tyr Ala Thr  
 210 215 220  
 His Arg Met Glu Glu Met Glu Glu Ala Phe Ala Ala Gly Asp Lys Asp  
 225 230 235 240  
 Ala Lys Leu Thr Pro Ala Val Lys Arg Asp Leu Arg Ala Ile Ala Arg  
 245 250 255  
 Asp Gly Arg Lys Ala Lys Asn His Leu Leu Glu Ala Asn Leu Arg Leu  
 260 265 270  
 Val Val Ser Leu Ala Lys Arg Tyr Thr Gly Arg Gly Met Ala Phe Leu  
 275 280 285  
 Asp Leu Ile Gln Glu Gly Asn Leu Gly Leu Ile Arg Ala Val Glu Lys  
 290 295 300  
 Phe Asp Tyr Ser Lys Gly Tyr Lys Phe Ser Thr Tyr Ala Thr Trp Trp  
 305 310 315 320  
 Ile Arg Gln Ala Ile Thr Arg Ala Met Ala Asp Gln Ala Arg Thr Ile  
 325 330 335  
 Arg Ile Pro Val His Met Val Glu Val Ile Asn Lys Leu Gly Arg Ile  
 340 345 350  
 Gln Arg Glu Leu Leu Gln Glu Leu Gly Arg Glu Pro Thr Pro Gln Glu  
 355 360 365  
 Leu Ser Lys Glu Met Asp Ile Ser Glu Glu Lys Val Leu Glu Ile Gln  
 370 375 380  
 Gln Tyr Ala Arg Glu Pro Ile Ser Leu Asp Gln Thr Ile Gly Asp Glu  
 385 390 395 400  
 Gly Asp Ser Gln Leu Gly Asp Phe Ile Glu Asp Ser Glu Val Val Val  
 405 410 415  
 Ala Val Asp Ala Val Ser Phe Thr Leu Leu Gln Asp Gln Leu Gln Asp  
 420 425 430  
 Val Leu Glu Thr Leu Ser Glu Arg Glu Ala Gly Val Val Lys Leu Arg  
 435 440 445  
 Phe Gly Leu Thr Asp Gly Met Pro Arg Thr Leu Asp Glu Ile Gly Gln  
 450 455 460

Val Tyr Gly Val Thr Arg Glu Arg Ile Arg Gln Ile Glu Ser Lys Thr  
465 470 475 480

Met Ser Lys Leu Arg His Pro Ser Arg Ser Gln Val Leu Arg Asp Tyr  
485 490 495

Leu Asp

<210> 5  
<211> 1892  
<212> DNA  
<213> *Corynebacterium glutamicum*

<220>  
<221> CDS  
<222> (226)..(1719)  
<223> sigA allele

<220>  
<221> mutation  
<222> (1466)..(1466)  
<223> Exchange of cytosine for thymine

<400> 5  
tgatcggtg accaactcta taagagatgc acctcaagtt tggggatact tattcggcgt 60  
ttctggggaa caaatagctt tccctattgt tgtatatagg tattegcact taagaaacat 120  
ctctcatgga aagaagctag gcggaaaggg cgttaagtac ttgccattta atcctcagca 180  
tcactcggat cagtcggaga tgtcgatgaa aatgcaccag gagcc gtg gag agc agc 237  
Val Glu Ser Ser  
1  
atg gta gaa aac aac gta gca aaa aag acg gtc gct aaa aag acc gca 285  
Met Val Glu Asn Asn Val Ala Lys Lys Thr Val Ala Lys Lys Thr Ala  
5 10 15 20  
cgc aag acc gca cgc aaa gca gcc ccg cgc gtg gca acc cca ttg gga 333  
Arg Lys Thr Ala Arg Lys Ala Ala Pro Arg Val Ala Thr Pro Leu Gly  
25 30 35  
gtc gca tct gag tct ccc att tcg gcc acc cct gcg cgc agc atc gat 381  
Val Ala Ser Glu Ser Pro Ile Ser Ala Thr Pro Ala Arg Ser Ile Asp  
40 45 50  
gga acc tca acc cct gtt gaa gct gct gac acc ata gag acc acc gcc 429  
Gly Thr Ser Thr Pro Val Glu Ala Ala Asp Thr Ile Glu Thr Thr Ala  
55 60 65  
cct gca gcg aag gct cct gcg gcc aag gct ccc gct aaa aag gtt gcc 477  
Pro Ala Ala Lys Ala Pro Ala Ala Lys Ala Pro Ala Lys Lys Val Ala  
70 75 80  
aag aag aca gct cgc aag gca cct gcg aaa aag act gtc gcc aag aaa 525  
Lys Lys Thr Ala Arg Lys Ala Pro Ala Lys Lys Thr Val Ala Lys Lys  
85 90 95 100



gcc aca acc gcc aag gct gca cct gca act gcc aag gac gaa aac gca Ala Thr Thr Ala Lys Ala Ala Pro Ala Thr Ala Lys Asp Glu Asn Ala 105 110 115	573
cct gtt gat gac gac gag gag aac ctc gct cag gat gaa cag gac ttc Pro Val Asp Asp Asp Glu Glu Asn Leu Ala Gln Asp Glu Gln Asp Phe 120 125 130	621
gac ggc gat gac ttc gta gac ggc atc gaa gac gaa gaa gat gaa gac Asp Gly Asp Asp Phe Val Asp Gly Ile Glu Asp Glu Glu Asp Glu Asp 135 140 145	669
ggc gtc gaa gcc ctc ggt gaa gaa agc gaa gac gac gaa gag gac ggc Gly Val Glu Ala Leu Gly Glu Glu Ser Glu Asp Asp Glu Glu Asp Gly 150 155 160	717
tca tcc gtt tgg gat gaa gac gaa tcc gca acc ctg cgt cag gca cgt Ser Ser Val Trp Asp Glu Asp Glu Ser Ala Thr Leu Arg Gln Ala Arg 165 170 175 180	765
aaa gat gcc gag ctc acc gct tcc gcc gac tct gtt cgc gct tac ctg Lys Asp Ala Glu Leu Thr Ala Ser Ala Asp Ser Val Arg Ala Tyr Leu 185 190 195	813
aag caa atc ggt aaa gtt gcc ctg ctg aac gct gaa cag gaa gtc tcc Lys Gln Ile Gly Lys Val Ala Leu Leu Asn Ala Glu Gln Glu Val Ser 200 205 210	861
ctg gca aag cgc atc gaa gca ggc ctt tac gcc acc cac cgc atg gag Leu Ala Lys Arg Ile Glu Ala Gly Leu Tyr Ala Thr His Arg Met Glu 215 220 225	909
gaa atg gaa gaa gct ttc gca gcc ggt gac aag gac gcg aaa ctc acc Glu Met Glu Glu Ala Phe Ala Ala Gly Asp Lys Asp Ala Lys Leu Thr 230 235 240	957
cca gcc gtc aag cgt gac ctc cgc gcc atc gct cgt gac ggc cgc aag Pro Ala Val Lys Arg Asp Leu Arg Ala Ile Ala Arg Asp Gly Arg Lys 245 250 255 260	1005
gcg aaa aac cac ctc ctg gaa gcc aac ctt cgt ctg gtt gtc tcc ctg Ala Lys Asn His Leu Leu Glu Ala Asn Leu Arg Leu Val Val Ser Leu 265 270 275	1053
gca aag cgc tac acc ggc cgt ggc atg gca ttc ctg gac ctc atc cag Ala Lys Arg Tyr Thr Gly Arg Gly Met Ala Phe Leu Asp Leu Ile Gln 280 285 290	1101
gaa ggc aac ctc ggt ctg att cgt gcc gta gag aag ttc gac tac tcc Glu Gly Asn Leu Gly Leu Ile Arg Ala Val Glu Lys Phe Asp Tyr Ser 295 300 305	1149
aag ggc tac aag ttc tcc acc tac gca acc tgg tgg atc cgt cag gca Lys Gly Tyr Lys Phe Ser Thr Tyr Ala Thr Trp Trp Ile Arg Gln Ala 310 315 320	1197
atc acc cgc gcc atg gcc gac caa gca cga acc atc cgt atc cca gtc Ile Thr Arg Ala Met Ala Asp Gln Ala Arg Thr Ile Arg Ile Pro Val 325 330 335 340	1245

cac atg gtt gaa gtg atc aac aaa ctt ggt cgc atc caa cgt gaa ctc 1293  
 His Met Val Glu Val Ile Asn Lys Leu Gly Arg Ile Gln Arg Glu Leu  
 345 350 355

ctt cag gaa ctc ggc cgc gaa cca acc cca cag gaa ctg tcc aaa gaa 1341  
 Leu Gln Glu Leu Gly Arg Glu Pro Thr Pro Gln Glu Leu Ser Lys Glu  
 360 365 370

atg gac atc tcc gag gaa aag gta ctg gaa atc cag cag tac gcc cgc 1389  
 Met Asp Ile Ser Glu Glu Lys Val Leu Glu Ile Gln Gln Tyr Ala Arg  
 375 380 385

gaa cca atc tcc ctg gac caa acc atc ggc gac gaa ggc gac agc cag 1437  
 Glu Pro Ile Ser Leu Asp Gln Thr Ile Gly Asp Glu Gly Asp Ser Gln  
 390 395 400

ctc ggc gac ttc atc gaa gac tcc gaa gtc gtc gtc gca gtc gac gcc 1485  
 Leu Gly Asp Phe Ile Glu Asp Ser Glu Val Val Val Ala Val Asp Ala  
 405 410 415 420

gtc tca ttc acc ctg ctg caa gac cag cta cag gac gtc cta gag acc 1533  
 Val Ser Phe Thr Leu Leu Gln Asp Gln Leu Gln Asp Val Leu Glu Thr  
 425 430 435

ctc tcc gaa cgt gaa gcc ggc gtg gtt aaa ctc cgc ttc gga ctc acc 1581  
 Leu Ser Glu Arg Glu Ala Gly Val Val Lys Leu Arg Phe Gly Leu Thr  
 440 445 450

gac gga atg cca cgc act tta gac gaa atc ggc caa gtt tac ggt gtc 1629  
 Asp Gly Met Pro Arg Thr Leu Asp Glu Ile Gly Gln Val Tyr Gly Val  
 455 460 465

acc cgt gag cgc atc cgc cag att gag tcc aag acc atg tct aag ctg 1677  
 Thr Arg Glu Arg Ile Arg Gln Ile Glu Ser Lys Thr Met Ser Lys Leu  
 470 475 480

cgc cac cca tca cgc tcc cag gtc ctt cgc gac tac ctg gac 1719  
 Arg His Pro Ser Arg Ser Gln Val Leu Arg Asp Tyr Leu Asp  
 485 490 495

taaaacccca gtcgggctca agaccggggcc gccactgttt tcctctgcgg ggaacggtgg 1779

tggcccgggtt tttctgttgc tttgggttcgg ctggtcacag ttcggctggg gtgttttaag 1839

tttgatttca cattgccgat ttctaaacgc cgagttctcg gattcgagac ctt 1892

<210> 6  
 <211> 498  
 <212> PRT  
 <213> Corynebacterium glutamicum

<400> 6  
 Val Glu Ser Ser Met Val Glu Asn Asn Val Ala Lys Lys Thr Val Ala  
 1 5 10 15  
 Lys Lys Thr Ala Arg Lys Thr Ala Arg Lys Ala Ala Pro Arg Val Ala  
 20 25 30  
 Thr Pro Leu Gly Val Ala Ser Glu Ser Pro Ile Ser Ala Thr Pro Ala  
 35 40 45

Arg Ser Ile Asp Gly Thr Ser Thr Pro Val Glu Ala Ala Asp Thr Ile  
 50 55 60  
 Glu Thr Thr Ala Pro Ala Ala Lys Ala Pro Ala Ala Lys Ala Pro Ala  
 65 70 75 80  
 Lys Lys Val Ala Lys Lys Thr Ala Arg Lys Ala Pro Ala Lys Lys Thr  
 85 90 95  
 Val Ala Lys Lys Ala Thr Thr Ala Lys Ala Ala Pro Ala Thr Ala Lys  
 100 105 110  
 Asp Glu Asn Ala Pro Val Asp Asp Asp Glu Glu Asn Leu Ala Gln Asp  
 115 120 125  
 Glu Gln Asp Phe Asp Gly Asp Asp Phe Val Asp Gly Ile Glu Asp Glu  
 130 135 140  
 Glu Asp Glu Asp Gly Val Glu Ala Leu Gly Glu Glu Ser Glu Asp Asp  
 145 150 155 160  
 Glu Glu Asp Gly Ser Ser Val Trp Asp Glu Asp Glu Ser Ala Thr Leu  
 165 170 175  
 Arg Gln Ala Arg Lys Asp Ala Glu Leu Thr Ala Ser Ala Asp Ser Val  
 180 185 190  
 Arg Ala Tyr Leu Lys Gln Ile Gly Lys Val Ala Leu Leu Asn Ala Glu  
 195 200 205  
 Gln Glu Val Ser Leu Ala Lys Arg Ile Glu Ala Gly Leu Tyr Ala Thr  
 210 215 220  
 His Arg Met Glu Glu Met Glu Glu Ala Phe Ala Ala Gly Asp Lys Asp  
 225 230 235 240  
 Ala Lys Leu Thr Pro Ala Val Lys Arg Asp Leu Arg Ala Ile Ala Arg  
 245 250 255  
 Asp Gly Arg Lys Ala Lys Asn His Leu Leu Glu Ala Asn Leu Arg Leu  
 260 265 270  
 Val Val Ser Leu Ala Lys Arg Tyr Thr Gly Arg Gly Met Ala Phe Leu  
 275 280 285  
 Asp Leu Ile Gln Glu Gly Asn Leu Gly Leu Ile Arg Ala Val Glu Lys  
 290 295 300  
 Phe Asp Tyr Ser Lys Gly Tyr Lys Phe Ser Thr Tyr Ala Thr Trp Trp  
 305 310 315 320  
 Ile Arg Gln Ala Ile Thr Arg Ala Met Ala Asp Gln Ala Arg Thr Ile  
 325 330 335  
 Arg Ile Pro Val His Met Val Glu Val Ile Asn Lys Leu Gly Arg Ile  
 340 345 350  
 Gln Arg Glu Leu Leu Gln Glu Leu Gly Arg Glu Pro Thr Pro Gln Glu  
 355 360 365  
 Leu Ser Lys Glu Met Asp Ile Ser Glu Glu Lys Val Leu Glu Ile Gln  
 370 375 380

Gln Tyr Ala Arg Glu Pro Ile Ser Leu Asp Gln Thr Ile Gly Asp Glu  
 385 390 395 400  
 Gly Asp Ser Gln Leu Gly Asp Phe Ile Glu Asp Ser Glu Val Val Val  
 405 410 415  
 Ala Val Asp Ala Val Ser Phe Thr Leu Leu Gln Asp Gln Leu Gln Asp  
 420 425 430  
 Val Leu Glu Thr Leu Ser Glu Arg Glu Ala Gly Val Val Lys Leu Arg  
 435 440 445  
 Phe Gly Leu Thr Asp Gly Met Pro Arg Thr, Leu Asp Glu Ile Gly Gln  
 450 455 460  
 Val Tyr Gly Val Thr Arg Glu Arg Ile Arg Gln Ile Glu Ser Lys Thr  
 465 470 475 480  
 Met Ser Lys Leu Arg His Pro Ser Arg Ser Gln Val Leu Arg Asp Tyr  
 485 490 495  
 Leu Asp

<210> 7  
 <211> 1689  
 <212> DNA  
 <213> Artificial sequence

<220>  
 <221> misc\_feature  
 <222> (1)..(1689)  
 <223> Description of the artificial sequence: PCR product containing an  
 n 3'-terminal region of the sigA allele (= sigA\_A414V allele) and  
 the downstream region

<220>  
 <221> mutation  
 <222> (854)..(854)  
 <223> Exchange of cytosine for thymine

<400> 7  
 acgaattccg acggcgatga cttcgtagac ggcacggaag acgaagaaga tgaagacggc 60  
 gtcgaagccc tcggtgaaga aagcgaagac gacgaagagg acggctcatc cgtttgggat 120  
 gaagacgaat ccgcaaccct gcgtcaggca cgtaaagatg ccgagctcac cgcttccgcc 180  
 gactctgttc gcgcttacct gaagcaaadc ggtaaagttg cctgctgaa cgctgaacag 240  
 gaagtctccc tggcaaagcg catcgaagca ggcctttacg ccacccaccg catggaggaa 300  
 atggaagaag ctttcgcagc cggtgacaag gacgcgaaac tcacccacgc cgtcaagcgt 360  
 gacctccgcg ccatcgctcg tgacggccgc aaggcgaaaa accacctoct ggaagccaac 420  
 cttcgtctgg ttgtctccct ggcaaagcgc tacaccggcc gtggcatggc attcctggac 480  
 ctcatccagg aaggcaacct cggcttgatt cgtgccgtag agaagttcga ctactccaag 540  
 ggctacaagt tctccaccta cgcaacctgg tggatccgtc aggcaatcac ccgcgccatg 600

```

gccgaccaag cacgaaccat ccgtatccca gtccacatgg ttgaagtgat caacaaactt 660
ggtcgcatcc aacgtgaact ccttcaggaa ctcggccgcg aaccaacccc acaggaactg 720
tccaaagaaa tggacatctc cgaggaaaag gtactggaaa tccagcagta cgcccgcgaa 780
ccaatctccc tggaccaaac catcgggcgcg gaaggcgaca gccagctcgg cgacttcac 840
gaagactccg aagtcgtcgt cgcagtcgcg gccgtctcat tcaccctgct gcaagaccag 900
ctacaggacg tcctagagac cctctccgaa cgtgaagccg gcgtgggtaa actccgcttc 960
ggactcaccg acggaatgcc acgcacttta gacgaaatcg gccaaagtta cgggtgcacc 1020
cgtgagcgca tccgccagat tgagtccaag accatgtcta agctgcgcca cccatcacgc 1080
tcccaggctc ttgcgcacta cctggactaa aaccccagtc gggctcaaga ccgggccgcc 1140
actgttttcc tctgcgggga acggtgggtg cccggttttt ctgttgcttt ggctcggtg 1200
gtcacagttc ggctggggtg ttttaagttt gatttcacat tgccgatttc taaacgccga 1260
gttctcggat tcgagacctt acctgcaatt tcacgggttac aatttctcgt tagcactttc 1320
gcgtactcaa tttcttatgt tcaatttcgg tcggaaaagt gccattttcc gacatcgctt 1380
tgagaaattg aatacaagaa attgagcgca aggtatcgaa cttgagaaat tgagcttttag 1440
cactcactcc cctgtttgat gaagtcagca aagccaaaag acgacttcac atctccgact 1500
tcttaacagc gacttctcgt ggctgacttc gctacctaaa cctgagtttc cttagcgaag 1560
tcgtgtgatg agaagtcgtg tgatgagaag ttgggtatga gaagtcagca cacgtgaagt 1620
cgcttttctt ccccggcacg ctcggagtag cgtgaaaggt ggaatggagc gaggtggaac 1680
ggaattcca 1689

```

```

<210> 8
<211> 20
<212> DNA
<213> Artificial sequence

```

```

<220>
<221> misc_feature
<222> (1)..(20)
<223> Description of the artificial sequence: primer sigA-1

```

```

<400> 8
tgatcggtg accaactcta

```

20

```

<210> 9
<211> 20
<212> DNA
<213> Artificial sequence

```

```

<220>
<221> misc_feature
<222> (1)..(20)
<223> Description of the artificial sequence: primer sigA-2

```

<400> 9  
aagggtctcga atccgagaac 20

<210> 10  
<211> 28  
<212> DNA  
<213> Artificial sequence

<220>  
<221> misc\_feature  
<222> (1)..(28)  
<223> Description of the artificial sequence: primer sigA\_XL-A1

<400> 10  
acgaattccg acggcgatga cttagtag 28

<210> 11  
<211> 28  
<212> DNA  
<213> Artificial sequence

<220>  
<221> misc\_feature  
<222> (1)..(28)  
<223> Description of the artificial sequence: primer sigA\_XL-A2

<400> 11  
tggaattccg ttccacctcg ctccattc 28

<210> 12  
<211> 20  
<212> DNA  
<213> Artificial sequence

<220>  
<221> misc\_feature  
<222> (1)..(20)  
<223> Description of the artificial sequence: primer sA\_1

<400> 12  
aagttctcca cctacgcaac  
20

International Application No.  
PCT/EP 02/11561

International Application No.  
PCT/EP 02/11561

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 C12N9/12 C12N15/31 C12N15/52 C07K14/34 C07K14/345  
C12N15/77 C12N1/21 C12P13/08

According to International Patent Classification (IPC) or to both national classification and IPC

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C12P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, WPI Data

### C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 1 108 790 A (KYOWA HAKKO KOGYO KK) 20 June 2001 (2001-06-20) ---	
A	WO 01 66573 A (BASF AG) 13 September 2001 (2001-09-13) ---	
A	HALGASOVA NORA ET AL: "Cloning and transcriptional characterization of two sigma factor genes, sigA and sigB, from <i>Brevibacterium flavum</i> ." CURRENT MICROBIOLOGY, vol. 43, no. 4, October 2001 (2001-10), pages 249-254, XP002227863 ISSN: 0343-8651 ----- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date of priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "A" document member of the same patent family

Date of the actual completion of the international search

20 January 2003

Date of mailing of the international search report

05/02/2003

**Name and mailing address of the ISA**  
European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer \_\_\_\_\_

Marinoni, J-C

# INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 02/11561

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>GOMEZ MANUEL ET AL: "sigA Is an essential gene in Mycobacterium smegmatis."  MOLECULAR MICROBIOLOGY,  vol. 29, no. 2, July 1998 (1998-07), pages  617-628, XP002227864  ISSN: 0950-382X</p>	
A	<p>OGUIZA J A ET AL: "Multiple sigma factor genes in brevibacterium lactofermentum: characterization of sigA and sigB"  JOURNAL OF BACTERIOLOGY, WASHINGTON, DC, US,  vol. 178, no. 2, January 1996 (1996-01),  pages 550-553, XP002183859  ISSN: 0021-9193</p>	



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 02/11561

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 1108790	A	20-06-2001	EP 1108790 A2	20-06-2001
			JP 2002191370 A	09-07-2002
			US 2002197605 A1	26-12-2002
WO 0166573	A	13-09-2001	AU 2390301 A	17-09-2001
			EP 1257649 A2	20-11-2002
			EP 1261718 A2	04-12-2002
			WO 0100843 A2	04-01-2001
			WO 0166573 A2	13-09-2001